

Statistical Mechanics of Charged Particles in Einstein-Maxwell-Scalar Theory

Kiyoshi Shiraishi*

Akita Junior College

Shimokitade-sakura, Akita-shi, Akita 010, Japan

Takuya Maki†

Kitasato University

Kitasato, Sagamihara-shi, Kanagawa 228, Japan

(February 3, 2008)

Abstract

We consider an N -body system of charged particle coupled to gravitational, electromagnetic, and scalar fields. The metric on moduli space for the system can be considered if a relation among the charges and mass is satisfied, which includes the BPS relation for monopoles and the extreme condition for charged dilatonic black holes. Using the metric on moduli space in the long distance approximation, we study the statistical mechanics of the charged particles at low velocities. The partition function is evaluated as the leading order of the large d expansion, where d is the spatial dimension of the system and will be substituted finally as $d = 3$.

*e-mail: g00345@sinet.ad.jp, shiraish@air.akita-u.ac.jp

†e-mail: maki@nls.kitasato-u.ac.jp

Solitons in field theories have attracted much attention in past two decades. The dynamics of solitons have been studied not only by numerical methods but also by analytical methods. If a multi-soliton solution saturates a certain Bogomol'nyi-type bound, there is no net static force among the solitonic objects described by such a solution. The existence of various types of forces is essential for the balanced situation. The low-energy interaction of such solitons can be described in terms of the space of parameters in static solutions (moduli space) [1].

Recently, the statistical mechanics of vortices in (2+1) dimensions was studied by Manton and Shah [2,3] under the assumption that the dynamics of vortices is well approximated by the geodesic motion on moduli space. In the present paper, we investigate thermodynamic behavior of the gas of the charged particles interacting by gravitational, electromagnetic, and scalar forces. This model can be regarded as a point particle limit of monopoles and/or black holes when there is a special relation among the charges and mass. We adopt the lagrangian of particles at low energies and at large separations to compute the partition function of the system.

Suppose the action for the N particles of mass m_α , electric charge q_α and scalar charge σ_α in the $(d+1)$ dimensional spacetime ($d \geq 3$) is given by

$$L = - \sum_{\alpha=1}^N \int ds_\alpha \left(m_\alpha - \sigma_\alpha \varphi - q_\alpha A_\mu \frac{\partial x_\alpha^\mu}{\partial s_\alpha} \right) + \int d^d x \frac{\sqrt{-g}}{16\pi} \left[\frac{1}{G} R - 2(\nabla\varphi)^2 - F^2 \right] + (\text{surface terms}) , \quad (1)$$

where R is the scalar curvature and φ is a real scalar field. The electromagnetic field A_μ is related to the field strength $F_{\mu\nu}$ by $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, where μ, ν run over $0, 1, \dots, d+1$. G is the Newton's constant. We will finally focus on the case of $d = 3$, but for later use in an approximation scheme, we leave d as an arbitrary integer for a while.

One of the present authors has obtained the low-energy lagrangian including Liénard-Wiechert potential in the large mutual distance limit [4,5]. Using abbreviated notations

$$e_{\alpha\beta}^2 \equiv q_\alpha q_\beta - \sigma_\alpha \sigma_\beta - \frac{2(d-2)}{d-1} G m_\alpha m_\beta , \quad (2)$$

$$\gamma_{\alpha\beta}^2 \equiv \left[\sigma_\alpha \sigma_\beta - \frac{2d}{d-1} G m_\alpha m_\beta \right] , \quad (3)$$

we can write the effective lagrangian for the particles as

$$\begin{aligned} L = & \frac{1}{2} \sum_{\alpha=1}^N m_\alpha \vec{v}_\alpha^2 \\ & - \sum_{\alpha < \beta} \frac{4\pi \gamma_{\alpha\beta}^2}{A_{d-1}} \frac{|\vec{v}_\alpha - \vec{v}_\beta|^2}{2(d-2) r_{\alpha\beta}^{d-2}} \\ & + \sum_{\alpha < \beta} \frac{4\pi e_{\alpha\beta}^2}{A_{d-1}} \frac{[\vec{v}_\alpha \cdot \vec{v}_\beta + (d-2)(\vec{n}_{\alpha\beta} \cdot \vec{v}_\alpha)(\vec{n}_{\alpha\beta} \cdot \vec{v}_\beta)]}{2(d-2) r_{\alpha\beta}^{d-2}} \\ & - \sum_{\alpha < \beta} \frac{4\pi e_{\alpha\beta}^2}{A_{d-1}} \frac{1}{(d-2) r_{\alpha\beta}^{d-2}} \\ & + \dots , \end{aligned} \quad (4)$$

where $A_{d-1} = \frac{2\pi^{d/2}}{\Gamma(d/2)}$,¹ $r_{\alpha\beta}$ is the distance between two point sources denoted by α and β , and $\vec{n}_{\alpha\beta}$ stands for the unit normal vector in the direction β - α . Here, the triple dots ($\dots$) contain $O(v^3)$ and $O(1/r^{2(d-2)})$ interactions. The radiation reaction is negligible at low velocities. Note that this lagrangian can be utilized for the system of magnetic monopoles, where the duality relation holds.

In eq. (4), the static interaction is cancelled out if

$$e_{\alpha\beta}^2 = 0 . \quad (5)$$

For the BPS monopoles in $(3+1)$ dimensions, if gravity is negligible ($G \rightarrow 0$), the relation $q_\alpha = \sigma_\alpha$ holds for each monopoles [6]; then the balance condition (5) is satisfied. For extreme dilatonic black holes [7,4,5],

$$\begin{aligned} q_\alpha &= \sqrt{\frac{2(d-2+a^2)}{d-1}} \sqrt{G} m_\alpha , \\ \sigma_\alpha &= a \sqrt{\frac{2}{d-1}} \sqrt{G} m_\alpha , \end{aligned} \quad (6)$$

where a is the dilaton coupling constant. The balance condition (5) is satisfied also in this case.

¹We use a different unit system from that in ref. [4,5] and fixed coefficients.

When the balance condition (5) is satisfied, there remains the interaction term which depends on the relative velocities. In this critical case, the slow motion of the well-separated particles is geodesic on the moduli space, and the evaluation of the partition function for the system of the particles reduces to the volume of the moduli space [2]. For non-critical cases, the deviation from the geodesic motion is expected to be modest if the couplings of the static potential and the other complicated velocity-dependent interaction are small [3]. Therefore we assume that we can express the partition function of the system in terms of a collective coordinates on the moduli space in the case of general charges and mass at least in the leading order calculation.

The lagrangian of the N -body system can be read as

$$L = \frac{1}{2} g_{AB} \dot{Q}^A \dot{Q}^B - V(Q) , \quad (7)$$

where $\{A\} = \{\alpha, i\}$ with $\alpha = 1, 2, \dots, N$ and $i = 1, 2, \dots, d$. Q is considered as the coordinates in Nd dimensional space. g_{AB} is called as the metric on moduli space of N -body system when the balance condition is satisfied. $V(Q)$ denotes the static potential.

The partition function is then given by

$$Z = \frac{1}{h^{Nd}} \int [dQ dP] e^{-\beta H(P, Q)} , \quad (8)$$

where $h = 2\pi\hbar$, $\beta = (k_B T)^{-1}$; T is the temperature of the system and k_B is the Boltzmann's constant.

The momentum corresponding to Q is defined as

$$P_A = \frac{\partial L}{\partial \dot{Q}^A} = g_{AB} \dot{Q}^B . \quad (9)$$

Using this canonical momentum, the Hamiltonian is expressed as

$$H = \frac{1}{2} g^{AB} P_A P_B + V(Q) , \quad (10)$$

where g^{AB} satisfies $g^{AC} g_{CB} = \delta_B^A$.

Performing the Gaussian integrals over P , we find

$$Z = \left(\frac{2\pi}{\beta h^2} \right)^{\frac{Nd}{2}} \int [dQ] \sqrt{\det g_{AB}(Q)} \exp[-\beta V(Q)] . \quad (11)$$

If the balance condition is satisfied, the partition function Z can be expressed by using the volume of the moduli space $\mathcal{V}_{\mathcal{M}}$ as

$$Z = \left(\frac{2\pi}{\beta h^2} \right)^{\frac{Nd}{2}} \mathcal{V}_{\mathcal{M}} , \quad (12)$$

where

$$\mathcal{V}_{\mathcal{M}} = \int [dQ] \sqrt{\det g_{AB}} . \quad (13)$$

We turn now to the evaluation of the partition function for the particle system governed by gravitational, electromagnetic, and scalar forces. Keeping terms of order v^2 and $1/r^{d-2}$, we recast the classical lagrangian for N identical particles of equal charges q , σ and mass m as

$$\begin{aligned} L = & \frac{1}{2} m \sum_{\alpha=1}^N \vec{v}_{\alpha}^2 \\ & - \frac{4\pi\gamma^2}{A_{d-1}} \sum_{\alpha<\beta} \frac{|\vec{v}_{\alpha} - \vec{v}_{\beta}|^2}{2(d-2) r_{\alpha\beta}^{d-2}} \\ & + \sum_{\alpha<\beta} \frac{4\pi e^2}{A_{d-1}} \frac{[\vec{v}_{\alpha} \cdot \vec{v}_{\beta} + (d-2)(\vec{n} \cdot \vec{v}_{\alpha})(\vec{n} \cdot \vec{v}_{\beta})]}{2(d-2) r_{\alpha\beta}^{d-2}} \\ & - \sum_{\alpha<\beta} \frac{4\pi e^2}{A_{d-1}} \frac{1}{(d-2) r_{\alpha\beta}^{d-2}} , \end{aligned} \quad (14)$$

where $\gamma^2 \equiv \sigma^2 - \frac{2d}{d-1} Gm^2$ and $e^2 \equiv q^2 - \sigma^2 - \frac{2(d-2)}{d-1} Gm^2$.

We use this lagrangian for well-separated particles to calculate the partition function. The approximation is justified if the average distance between particles is large. Thus we have to consider a dilute gas of the particles in our analysis.

From the lagrangian (14), we obtain the metric g_{AB} as follows:

$$g_{AB} = m I_{AB} - \gamma^2 U_{AB} + e^2 V_{AB} , \quad (15)$$

with

$$I_{AB} = \delta_{ij} \otimes \delta_{\alpha\beta} , \quad (16)$$

$$U_{AB} = \delta_{ij} \otimes u_{\alpha\beta} , \quad (17)$$

where

$$u = \begin{pmatrix} \sum_{\alpha \neq 1}^N \phi_{1\alpha} & -\phi_{12} & -\phi_{13} & \cdots & -\phi_{1N} \\ -\phi_{12} & \sum_{\alpha \neq 2}^N \phi_{2\alpha} & -\phi_{23} & \cdots & -\phi_{2N} \\ -\phi_{13} & -\phi_{23} & \sum_{\alpha \neq 3}^N \phi_{3\alpha} & \cdots & -\phi_{3N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\phi_{1N} & -\phi_{2N} & -\phi_{3N} & \cdots & \sum_{\alpha \neq N}^N \phi_{N\alpha} \end{pmatrix} , \quad (18)$$

$$V_{AB} = \begin{cases} (\delta_{ij} + n_{\alpha\beta,i} n_{\alpha\beta,j}) \phi_{\alpha\beta} & (\alpha \neq \beta) \\ 0 & (\alpha = \beta) \end{cases} , \quad (19)$$

and

$$\phi_{\alpha\beta} = \frac{4\pi}{A_{d-1}(d-2) r_{\alpha\beta}^{d-2}} , \quad (20)$$

First we consider the case that $e^2 = 0$. Then we get

$$\begin{aligned} \sqrt{\det g_{AB}} &= m^{\frac{Nd}{2}} \exp \frac{1}{2} \text{Tr} \ln(I_{AB} - \frac{\gamma^2}{m} U_{AB}) \\ &= m^{\frac{Nd}{2}} \exp \frac{d}{2} \text{Tr} \ln(\delta_{\alpha\beta} - \frac{\gamma^2}{m} u_{\alpha\beta}) \\ &= m^{\frac{Nd}{2}} \exp \left\{ -\frac{d}{2} \left[\frac{\gamma^2}{m} \text{Tr} u + \frac{1}{2} \left(\frac{\gamma^2}{m} \right)^2 \text{Tr} u^2 + \cdots \right] \right\} . \end{aligned} \quad (21)$$

Taking the limit $d \rightarrow \infty$, $\frac{d}{m} \equiv \frac{1}{\tilde{m}}$ fixed,² one finds that $\frac{1}{\tilde{m}^2} \phi^2$ contribution from $\text{Tr} u^2$ is suppressed by a factor $1/d$ in comparison with that from $\text{Tr} u$ in the expression (21), and also higher order terms of $\frac{1}{\tilde{m}} \phi$ from $\text{Tr} u^n$ ($n \geq 2$) or the product of them are more or so suppressed. Thus, according to this “large d ” approximation, the expression (21) reduces to

²At the same time, we have to set $Gm^2 = \tilde{G}\tilde{m}^2$.

$$\begin{aligned}
\sqrt{\det g_{AB}} &\approx m^{\frac{Nd}{2}} \exp \left[-\frac{\gamma^2 d}{2m} \text{Tr } u \right] \\
&= m^{\frac{Nd}{2}} \exp \left(-\frac{\gamma^2 d}{m} \sum_{\alpha < \beta} \phi_{\alpha\beta} \right) \\
&= m^{\frac{Nd}{2}} \exp \left(-\frac{\gamma^2 d}{m} \sum_{a=1}^M \phi_a \right), \tag{22}
\end{aligned}$$

where a represents the pair of indices $\{\alpha, \beta\}$, such as $\{1, 2\}$, and $M = N(N-1)/2$.

In this leading $1/d$ approximation scheme, the contribution of V_{AB} is neglected even for finite e^2 , since V_{AB} is traceless.

One can find that this closely resemble the contribution of static potential to the partition function,³ which can be written by

$$\exp(-\beta V) = \exp \left(-e^2 \beta \sum_{a=1}^M \phi_a \right). \tag{23}$$

Thus the integral over Q in eq. (11) can be written in a combined form

$$\frac{V^N}{N!} m^{\frac{Nd}{2}} e^{NW}, \tag{24}$$

where V is the volume of the system. As the thermodynamic limit $N \rightarrow \infty$ and $V \rightarrow \infty$ is taken, W is given by [8]

$$NW = \frac{1}{V^N} \sum_{n_1, \dots, n_M}' \left(-\frac{\gamma^2 d}{m} - e^2 \beta \right)^{\sum_a^M n_a} \frac{\int (\phi_1^{n_1} \phi_2^{n_2} \dots \phi_M^{n_M})_{irr} \prod_{\alpha=1}^N d^d x_\alpha}{n_1! n_2! \dots n_M!}, \tag{25}$$

where *irr* stands for the irreducible set and $\sum'$ means the sum excluding the term of $n_1 = n_2 = \dots = n_M = 0$.

Here we will use the ring approximation [8]; we pick up the irreducible set of a ring shape. Using this approximation with the Fourier transform, we get

$$\begin{aligned}
W_0 &= \frac{1}{2V} \sum_{n=2}^{\infty} \frac{(-\gamma^2 d - e^2 \beta m)^n \rho^{n-1}}{m^n n} \sum_{\vec{k}} \left(\frac{4\pi}{|\vec{k}|^2} \right)^n \\
&= \frac{1}{2\rho V} \sum_{\vec{k}} \left\{ -\ln \left[1 + \rho \frac{\kappa^2}{|\vec{k}|^2} \right] + \rho \frac{\kappa^2}{|\vec{k}|^2} \right\}, \tag{26}
\end{aligned}$$

³Thus positive γ^2 give rise to an effective repulsive force between the particles.

where the density of particles is defined as $\rho = N/V$ and $\kappa^2 = 4\pi \left(\frac{\gamma^2 d}{m} + e^2 \beta \right) \rho$. Here the suffix 0 means that we used the ring approximation.

Consequently, we have the result for $d = 3$, though d may not be very big:

$$\begin{aligned} Z &\approx \frac{V^N}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3N}{2}} e^{NW_0} \\ &= \frac{V^N}{N!} \left(\frac{2\pi m}{\beta h^2} \right)^{\frac{3N}{2}} \exp \left\{ \frac{N}{12\pi} \left[\frac{4\pi}{m} (3\gamma^2 + e^2 \beta m) \right]^{\frac{3}{2}} \rho^{\frac{1}{2}} \right\}. \end{aligned} \quad (27)$$

This yields the equation of state given by

$$\frac{pV}{Nk_B T} = 1 - \frac{1}{24\pi} \left[\frac{4\pi}{m} (3\gamma^2 + e^2 \beta m) \right]^{\frac{3}{2}} \rho^{\frac{1}{2}}, \quad (28)$$

where p is the pressure.

The deviation from an ideal gas is independent of the temperature for $e^2 = 0$. This is true if one may use any approximation, because the partition function is proportional to the volume of the moduli space $\mathcal{V}_{\mathcal{M}}$ in this special case. One can also see from eq. (28) that a possible phase transition occurs at high density, but the large distance approximation may become unreliable in such a case.

More interesting aspect is the appearance of the instability of the system. In the case of $\kappa^2 < 0$, imaginary contributions appear in the free energy ($\propto \ln Z$), which indicates the instability or the absence of the thermodynamical equilibrium in the system. The “phase” diagram for charges is shown in FIG. 1, where the possible sign of κ^2 is exhibited. When the charges fall into the upper-right region, the system is stable for any temperature. In the lower-left region, the system is always unstable. In the upper-left region, the system becomes unstable at low temperature. In the lower-right region, the system becomes unstable at high temperature.

For charged dilaton black holes [7] with $a^2 < 3$, the allowed charges fall into the region that $\kappa^2 < 0$ for any value of βm in the diagram. The instability may be considered as a characteristic feature of gravitating systems. For $a^2 > 3$, if $\gamma^2 > 0$ is satisfied, κ^2 becomes

positive when the temperature is sufficiently high.⁴

The charges of the *extreme* dilaton black holes correspond to the line described by $q^2 = \sigma^2 + Gm^2$. The system of the extreme black holes is always unstable for $a^2 < 3$, while for $a^2 > 3$ the system is stable according to the present analysis. The extreme case for $a^2 = 3$ corresponds to the point K in FIG. 1. This is also corresponding to the Kaluza-Klein monopole [9]. In this case, since there is no interaction of the order of v^2 , the system is equivalent to the classical ideal gas of free particles. On the other hand, the extreme Reissner-Nordström black hole corresponds to the point E [10]. In this case, the system is essentially unstable.

One must also bear in mind that the present estimation of the partition function has been obtained by using several approximations. More precise approximation schemes should be studied for some specific models of extended objects as well as of the point particles. In the case that the minute structure of moduli space is well-known, such as for the BPS monopoles [1,11] and charged black holes [12,5,13], application of the moduli metric to the thermodynamics requires further work with and without quantum mechanical considerations.

Finally, we expect that the result of the work by Gibbons and Manton [6] on the moduli space of well-separated dyons may be utilized to study the thermodynamics of dyons. Further investigation will be reported elsewhere.

⁴Actually, the charges of the any dilatonic black holes fall into the region between the lines $q^2 = \sigma^2 + Gm^2$ and $q^2 = \sigma^2$.

REFERENCES

- [1] N. Manton, Phys. Lett. **B110**, 54 (1982).
R. S. Ward, Phys. Lett. **B158**, 424 (1985).
M. F. Atiyah and N. J. Hitchin, *The Geometry and Dynamics of Magnetic Monopoles* (Princeton University Press, Princeton, 1988).
- [2] N. S. Manton, Nucl. Phys. **B400**, 624 (1993).
P. A. Shah and N. S. Manton, J. Math. Phys. **35**, 1171 (1994).
- [3] P. A. Shah, preprint DAMTP 94-86.
- [4] K. Shiraishi, J. Math. Phys. **34**, 1480 (1993).
- [5] K. Shiraishi, Nucl. Phys. **B402**, 399 (1993).
- [6] N. S. Manton, Phys. Lett. **B154**, 397 (1985); (E) **B157**, 475 (1985).
G. W. Gibbons and N. Manton, Phys. Lett. **B355**, 32 (1995).
- [7] G. W. Gibbons and K. Maeda, Nucl. Phys. **B298**, 741 (1988).
D. Garfinkle, G. Horowitz and A. Strominger, Phys. Rev. **D43**, 3140 (1991); (E) **D45**, 3888 (1992).
- [8] R. Abe, *Toukeirikigaku (Statistical mechanics)* (Tokyo University Press) (in Japanese).
- [9] P. J. Ruback, Commun. Math. Phys. **107**, 93 (1986).
- [10] G. W. Gibbons and P. J. Ruback, Phys. Rev. Lett. **57**, 1492 (1986).
- [11] G. W. Gibbons and N. S. Manton, Nucl. Phys. **B274**, 183 (1986).
B. J. Schroers, Nucl. Phys. **B367**, 177 (1991).
- [12] R. C. Ferrell and D. M. Eardley, Phys. Rev. Lett. **59**, 1617 (1987).
J. Traschen and R. Ferrell, Phys. Rev. **D45**, 2628 (1992).
- [13] K. Shiraishi, Int. J. Mod. Phys. **D2**, 59 (1993); preprint AJC-HEP-26, gr-qc/9507029.

FIGURES




FIG. 1. The phase diagram for charges of the particles, for $d = 3$.